

ORAL PATHOLOGICAL DIAGNOSTIC

THE COUNCIL FOR TOBACCO RESEARCH - U.S.A.

SUCCESSOR TO THE  
TOBACCO INDUSTRY RESEARCH COMMITTEE

635 THIRD AVENUE  
NEW YORK, N. Y. 10017

COMMITTEE:

Dr. Reimann, Chm.  
Dr. Andervont  
Dr. Little

Application For Research Grant

Date: 31 January 1966

1. Name of Investigator: John H. Manhold, Jr., D.M.D., M.A., F.A.C.D., F.I.C.D.
2. Title: Professor and Director of Pathology and Oral Diagnosis for the College of Dentistry
3. Institution & Address: New Jersey College of Medicine and Dentistry  
Jersey City, New Jersey 07304
4. Project or Subject: To study the purported relationship between smoking and changes in human oral tissue in vivo by routine microscopy, differential staining, and micro-respirometer methods, and to further examine statistically two series of data presently in the principal investigator's possession.

5. Detailed Plan of Procedure (Use additional pages if more space is required.)

Introductory data -- The relationship between smoking and malignant or premalignant changes in bronchial and lung tissue has been much studied. Additionally, oesophageal changes with smoking have been studied at some length by, among others Wynder and Gross<sup>1</sup> and Auerbach<sup>2</sup> (who incidentally also is one of the investigators who has published several articles on bronchial and lung changes associated with smoking).<sup>3-7</sup> The oral cavity per se has not been as well studied although many statistics and much conjecture exist in regard to a purported relationship between smoking and the occurrence of malignancy in this area. However, the statistics stem largely from linear correlations which continue to offer considerable controversy and the conjecture, along with some of the statistics, stem from animal experimentation done with the use of "smoke boxes" and/or the application of various carcinogenic agents. Actual human in vivo experimentation from other than the statistical viewpoint, is not extensive. Kirsch<sup>8</sup> for example, has suggested the existence of a difference in salivary pH between smokers and non-smokers. Pindborg<sup>9</sup> from the results of microscopic study of a small sample, has suggested that cigarette smoking produces a hyper-parakeratotic condition in the oral cavity. Wrubel and Scopp<sup>10</sup> using oral exfoliative cytologic methods have reported no change in the hard palate and buccal mucosa with cessation of smoking. On the other hand, Zimmerman and Zimmerman<sup>11</sup> using the same method, have reported significant differences in oral tissue keratinization between smokers and non-smokers.

A certain lack of confidence exists among many pathologists in regard to oral exfoliative cytologic techniques. Peters<sup>12</sup> has demonstrated some basis for this lack of confidence, and Manhold et al.<sup>13</sup> have demonstrated a further apparent unreliability of the oral cytologic technique by not being able to find histological evidence to corroborate the cytologic results reported by Derbyshire and Manhold.<sup>14</sup>

1005101385

Nevertheless, these various studies by Kirsch, Pindborg, Wrubel and Scopp and Zimmerman and Zimmerman, when considered in relation to studies done by Cahn and associates,<sup>15, 16, 17</sup> suggest the possibility of more exact human *in vivo* investigation of the oral tissue-smoking problem. For example, Cahn, Eisenberg and Blake suggested in 1961 that it may be possible to distinguish potentially malignant epithelium from benign epithelium by study of the basement membrane with Periodic Acid Schiff Stain and in 1962 these same authors suggested that the existence of parakeratosis without glycogen may be "a sinister sign in the prognosis of white lesions of the mouth."<sup>16</sup> They also suggested in this same study that a difference in RNA could be seen with methyl-green pyronine.

It is proposed, therefore, to study the purported relationship between smoking and changes in human oral tissue *in vivo* by routine microscopy, differential staining, and microrespirometer methods, and to further examine statistically two series of data presently in the principal investigator's possession in the following manner:

#### PROCEDURE

##### Part I

It is proposed that a group of individuals who are heavy cigarette smokers and a group who are non-smokers have an area biopsied from the oral mucosa and that this biopsy be submitted to histologic study with H & E, Periodic Acid Schiff, Methyl-green Pyronine, and Feulgen Stain methods. Simultaneously part of these tissues will be subjected to tissue microrespirometry to ascertain whether any oxygen consumption difference appears to exist between the tissue of the smokers and non-smokers.

The staining techniques employed will be standard and will be utilized to compare with the results presented by Pindborg, Wrubel and Scopp, Zimmerman and Zimmerman and Cahn et al. The microrespirometer technique to be utilized here is fundamentally that initiated by Kirk.<sup>18</sup> It will employ the Stern-Kirk microrespirometer in the manner described by Manhold and associates,<sup>19, 20</sup> will allow comparisons with the values reported by these authors for normal and inflamed tissues.

##### Part II

In addition to this procedure the following is suggested:

Over the past seven years the proposed principal investigator has conducted a Biopsy Service for dentists of the State of New Jersey and environs. This service now processes in excess of 1,000 specimens per year. Over the past year and one-half there have been, among the biopsies submitted, 37 patients whom we have diagnosed as having lesions of a fundamentally hyperkeratotic nature; 8 patients who have been diagnosed as having invasive carcinoma; and there have been 2 patients who have been diagnosed as having leukoplakia. I propose first to contact the surgeons who submitted these specimens so as to gain pertinent data in regard to smoking habits, use of alcohol, and any medical or oral history which could be pertinent.

With these data for comparison, sections from blocks of these lesions will be cut and re-examined microscopically with appropriate H & E and differential staining as described above in Part I. (If any significant trend appears from this preliminary perusal, additional information could later be gained by tracing all cases of this type back through the remaining six years of our Biopsy Service. This work could also then be expanded to study any epithelial hyperplastic change. Further work in this area would, as stated, depend upon the apparent fruitfulness of this preliminary survey and would however, constitute another area of endeavor, which would necessitate further planning and additional funds.)

##### Part III

Because production of oral carcinoma is thought by many to stem largely from irritation, and because changes in the normal physiology of components of the oral cavity can be thought to be part of this irritational factor and also because Kirsch<sup>8</sup>

1005101386

has suggested that a difference exists in the pH of the saliva of smokers and non-smokers, it might well prove interesting to ascertain whether this reported difference can be replicated in another sample. We have already accumulated data on a number of individuals and established a relationship between saliva, dental calculus formation and Blood Serum Phosphorus.<sup>27</sup> Thus we have pH and quantity of saliva on this group. We have, additionally but as yet unanalyzed, data on these individuals as to whether or not they are smokers. To examine these relationships will require merely a T-test or critical ratio on these data.

1005101387

BIBLIOGRAPHY

1. Wynder, E. L. and Bross, I. J.: Study of Etiological Factors in Cancer of the Oesophagus. Cancer. 14: 389, 1961.
2. Auerbach, O., Stout, A. P., Hammond, E. C., and Garfinkel, L.: Histologic Changes in Oesophagus in Relation to Smoking Habits. Archives of Environmental Health. 11: 4, 1965.
3. Ibid: Changes in Bronchial Epithelium in Relation to Cigarette Smoking and in Relation to Lung Cancer. New England J. Med. 265: 253, 1961.
4. Ibid: Changes in Bronchial Epithelium in Relation to Sex, Age, Residence, Smoking and Pneumonia. New England J. Med. 267: 111, 1962.
5. Ibid: Bronchial Epithelium in Former Smokers. New England J. Med. 267: 119, 1962.
6. Ibid: Smoking Habits and Age in Relation to Pulmonary Changes. New England J. Med. 269: 1045, 1963.
7. Ibid: Inter-relationship Among Various Histologic Changes in Bronchial Tubes and in Lung Parenchyma. Am. Rev. Resp. Dis. 90: 857, 1964.
8. Kirsch, Th.: Universitätsklinik und Poliklinik für Mund-Zahn- und Kieferkrankheiten, Heidelberg, Germany. Über den pH-Wert des menschlichen Speichels. Dent. Zahnärztl. Arch. 18: 174, 1963.
9. Pindborg, J. H.: Tobacco Induced Leukoplakias. Acta. Odont. Scandinavica 22: 499, 1964.
10. Wrubel, G. J. and Scopp, I. W.: A Study of the Exfoliative Cytology of the Hard Palate and Buccal Mucosa Following Cessation of Smoking. J. Dent. Res. 40: 341, 1961.
11. Zimmerman, E. R. and Zimmerman, A. L.: Effects of Race, Age, Smoking Habits, Oral and Systemic Disease on Oral Exfoliative Cytology. J. Dent. Res. 44: 627, 1965.
12. Peters, Hannah: Cytologic Smears from the mouth. Am. Jour. Clin. Path. 29: 219, 1958.
13. Manhold, J. H., Franzetti, J. and Fitzsimmons, L. J.: Effect of the Electric Toothbrush on Human Gingiva: Histologic and Microrespirometer Evaluation. Jour. Periodont. 36: 135, 1965.
14. Derbyshire, J. C. and Mankodi, S. M.: Gingival Keratinization with Hand and Electric Toothbrushes: A Cytologic Comparison. J. Am. Dent. A. 68: 255, 1964.

1005101388

15. Cahn, L., Eisenberg, L. and Blake, M. N.: Histochemical Analysis of White Lesions of the Mouth: I. The Basement Membrane. Jour. Oral Surg., Oral Med., Oral Path. 14: 596, 1961.
16. Cahn, L. R., Eisenberg, L. and Blake, M. N.: Histochemical Analysis of White Lesions of the Mouth: II. Analysis of Glycogen Content. Jour. Oral Surg., Oral Med., Oral Path. 15: 458, 1962.
17. Cahn, L. R.: The Early Detection of Cancer of the Mouth. Brit. Dent. J. 111: 285, 1961.
18. Kirk, Paul L.: Quantitative Ultramicroanalysis. John Wiley & Sons, N. Y. 1950. pg. 235 ff.
19. Manhold, J. H., Bolden, T. E., and Katz, S.: Microrespirometer Technique for Study of Human Gingival Tissue: Preliminary Report. Jour. Dent. Research 39: 746, 1960.
20. Manhold, J. H., Bolden, T. E., and Hall, M. L.: Preliminary Report on Respiration Changes in Rat gland Tissue as a Result of Ligation. Abstract. Jour. Dent. Research 40: 647, 1961.
21. Manhold, J. H., and Bolden, T. E.: Investigation of the Tissue Oxygenating Properties of Proprietary Substances by use of a Microrespirometer Technique: I. In Vitro Study of Kasdenol. Annals of Dentistry, December 1961.
22. Volpe, A. R., Manhold, J. H.: Microrespirometer Technique for Study of Human Gingival Tissue. Jour. Dent. Research 41: 420, 1962.
23. Volpe, A. R., Manhold, J. H., and Manhold, B. S.: Effect of Age and Other Factors Upon Normal Gingival Tissue Respiration. Jour. Dent. Research 41: 1060, 1962.
24. Manhold, J. H., and Volpe, A. R.: Effect of Inflammation in the Absence of Proliferation on  $O_2$  Consumption of Gingival Tissue. Jour. Dent. Research 42: 103, 1963.
25. Manhold, J. H., Volpe, A. R., and DeCroce, R.: Relationship Between Salivary Respiration and Periodontal Disease: A Preliminary Report. Preprinted abstracts of the I.A.D.R. Meeting of March 1963.
26. Manhold, J. H., Manhold, B. S., and Volpe, A. R.: Relation Between Salivary Calculus and Blood Serum Phosphorus. Abstract, Jour. Dent. Research 43: 838, 1964.
27. Manhold, J. H., Volpe, A. R., and Manhold, B. S.: Interrelationship between Salivary Calculus Formation, Blood Serum Phosphorus Level, and Salivary Respiration ( $QO_2$ ). Jour. Perio. 35: 471, 1965.

1005101389

JOHN H. MANHOLD, JR., B.A., D.M.D., M.A., F.A.C.D., F.I.C.D.

**REDACTED**

2. Professional History in Brief:

**REDACTED**

3. Education:

B.A., University of Rochester 1940  
D.M.D., Harvard University 1944  
Fellow in Pathology, University of Rochester 1944  
Certificate in Medical Aspects of Atomic, Bacteriological, Chemical  
Warfare: Army Chemical Corps School, Maryland 1950  
M.A., in Psychology, Washington University 1956

4. Honors:

International College of Dentists (Fellow)  
American College of Dentists (Fellow)  
Society of the Sigma Xi (Member)  
Omicron Kappa Upsilon (Member)

5. Biography in Who's Who in the East (since 1950), American Men of Science (Vol. 11), and Leaders in American Science.

6.

b.

**REDACTED**

c.

7. Positions (from present to earliest)

Professor and Director of Pathology and Oral Diagnosis, Seton Hall  
(now New Jersey) College of Medicine and Dentistry, 1957-;  
Secretary to the Executive Faculty, 1957-; Coordinator of  
Cancer Teaching, 1956-; Attending in Pathology, Jersey City  
Medical Center 1952-; Coordinator of Research 1956-1960.  
Associate Professor and Director of Pathology, College of Dentistry,  
Seton Hall College of Medicine and Dentistry, 1957-1957.  
Assistant Professor and Director of Pathology, College of Dentistry,  
Seton Hall College of Medicine and Dentistry, 1956-1957  
(Interim Appointment).

1005101390

Assistant Professor and Acting Head of the Departments of General and Oral Pathology, Washington University School of Dentistry 1954-1956.

Research Officer and for several months Senior Research Officer Present, Aviation Psychology Laboratory and Research Pathologist, Naval School of Aviation Medicine, N.A.S., Pensacola, Florida 1952-1954.

Assistant Force Dental Officer, Fleet Marine Forces of the Pacific, 1951-1952.

Assistant Research Officer and Lecturer on Medical Aspects of Atomic Explosion, U.S.N.T.C., Great Lakes, III., 1950-1951.

Director, Cancer Teaching Program and Instructor, Oral Pathology, Tufts University College of Dentistry 1948-1950.

Private practice for a short time after World War II in which service was performed in the Navy for two years.

Designated by the Bureau of Medicine and Surgery of the United States Navy as "Qualified for Independent Research Duty".

Formerly Principle Investigator of U.S. Navy Research Project NM 001 057.11

#### 8. Publications:

##### A. Books

- 1) INTRODUCTORY PSYCHOSOMATIC DENTISTRY by Manhold, J. H., Jr. Appleton-Century-Crofts, New York, 1956.
- 2) OUTLINE OF PATHOLOGY by Manhold, J. H., and Bolden, T. E. W. B. Saunders Co., Philadelphia, 1960.
- 3) Dental Clinics of North America, Contributing Author, W. B. Saunders Co., Philadelphia, November 1962, Chapter on "The Psychosomatic Process in Dental Disease".
- 4) Orban's Periodontia, Ed. 2, Contributing Author, C. V. Mosby Co., St. Louis, 1963.
- 5) Pharmacotherapeutics in Oral Disease, Contributing Author, McGraw-Hill Co., New York, 1964, Section on "Traumatic Lesions of the Mouth".
- 6) CLINICAL ORAL DIAGNOSIS by Manhold, J. H., Jr., McGraw-Hill, New York, 1965.

##### b. Papers (Abstracts are not included unless subsequently published as complete papers on the subject presenting substantially additional material.)

##### Early Research Reports:

- 1) Manhold, J. H., and Manhold, B. S.: A Preliminary Report on the Study of the Relationship of Psychosomatics to oral Conditions - Relationship of Personality to Dental Caries. Science 110:585, 1949.
- 2) Glickman, I., Turesky, S., and Manhold, J.: The Oxygen Consumption of Healing Gingiva. Jour. Dent. Research 29: 429, 1950.

1005101391

- 3) Manhold, J.: Report of a Study of Relationship of Personality Variables to Periodontal Conditions. Jour. Periodontology, 25:248, 1953.
- 4) Manhold, J., and Rosenberg, N.: Study of the Possible Relationship of Personality Variables to Dental Cavities. Jour. Dent. Research 33:357, 1954.
- 5) Manhold, J., and Izard, C.: The Relationship of Dental Caries to General Health. Science, 120:892, 1954.
- 6) Manhold, J. H., and Jones, M. B.: A Study of Social Attitudes In Relation to Dental Caries Formation. Washington Univ. Dental Jour. XXII, #2, May 1956. (Recalled after acceptance by Jour. Dent. Research because material covered in book INTRODUCTORY PSYCHOSOMATIC DENTISTRY which was in press and to be published before Jour. Dent. Research paper would go to press.)

U.S. Navy Research Reports Published (1950-1955):

- 7) Manhold, J.: A Preliminary Report on the Study of the Relationship of Psychosomatics to Oral Conditions II. Relationship of Personality to Periodontal Conditions. Great Lakes, XII., 1950.
  - 8) Izard, C., and Manhold, J.: Correlation of Peer Leadership Ratings; I. Medical Complaints, Nav. Sch. Av. Med., Pensacola, Florida 27 April 1954.
  - 9) Manhold, J.: Re-evaluation Study of the Dental Complaints of Flying Personnel. Special Research Report #54-1, Nav. Sch. Av. Med., N.A.S., Pensacola, Florida 11 January 1954.
  - 10) Jones, M. B., and Manhold, J. H.: Authoritarianism and Physical Fitness, Nav. Sch. Av. Med., Pensacola, Florida, 15 March 1953.
  - 11) Manhold, J. H., and Jones, M. B.: Authoritarianism and Dental Caries, Nav. Sch. Av. Med., Pensacola, Florida, 15 March 1955.
- Additional Naval Research Reports Published (under Navy Research Project NM 001 057.11).

- 12) NM 001 057.11.01 Further Studies of the Relationship of Personality Variables to Dental Caries, Nav. Sch. Av. Med., Pensacola, Florida, 7 May 1953.
- 13) NM 001 057.11.02 Dental Complaints of Flying Personnel, Nav. Sch. Av. Med., Pensacola, Florida, 25 November 1953.
- 13) NM 001 057.11.03 A Study of Psychosomatic Factors in Oral Pathology, Nav. Sch. Av. Med., Pensacola, Florida, 12 April 1954.
- 15) NM 001 057.11.04 The Relationship of Dental Caries to General Health, Nav. Sch. Av. Med., Pensacola, Florida, 8 June 1954.
- 16) NM 001 057.11.05 A Comparative Study of Several Methods of Measuring Dental Caries in Man. Nav. Sch. Av. Med., Pensacola, Florida, 2 September 1954.

Papers since affiliating with Seton Hall:

- 17) Manhold, J. H.: Biopsy from the Pathologist's Viewpoint. N. Y. Jour. Dent. XXVII, #5, May 1957, pg. 193 ff.
- 18) Manhold, J. H.: Effects of Radiation. N. Y. State Dent. Jour. 24:117, 1958.

1005101392



- 19) Manhold, J. H., and Hafner, A. J.: Dental Caries and Psychological Factors. Jour. Clinical Psychol. XIV, #3, 319, 195C.
- 20) Manhold, J. H.: The Role of Psychologic Predisposition in Dental Cavity Formation. Jour. Dent. Research 37:330, 195C.
- 21) Manhold, J. H.: Preliminary Interrelational Study of Dental Caries in 1000 Adult Patients. Abstract. Jour. Dent. Research 37:77, 195C.
- 22) Manhold, J. H.: A Pilot Experiment Employing Animals for the Study of Dental Psychosomatic Tenets. N. Y. Jour. Dent. 2C:343, 1958.
- 23) Bolden, T. E., Manhold, J. H., Jr., and Barnes, S. M.: The Effect of Age on the Exorbital Lacrimal Gland of the Rat. Quarterly, Nat'l. Dent. Assoc., 1C:105, 1960.
- 24) Manhold, J. H., Bolden, T. E., and Katz, S.: Microrespirometer Technique for Study of Human Gingival Tissue; Preliminary Report. Jour. Dent. Research 39:746, 1960.
- 25) Manhold, J. H., Bolden, T. E., and Hall, M. L.: Preliminary Report on Respiration Changes in Rat Gland Tissue as a Result of Ligation. Abstract. Jour. Dent. Research 40: 647, 1961.
- 26) Bolden, T. E., and Manhold, J. H.: Comparative Study of Submaxillary and Lacrimal Glands of the Rat. Abstract. Jour. Dent. Research 40:690, 1961.
- 27) Manhold, J. H., and Bolden, T. E.: Investigation of the Tissue Oxygenating Properties of Proprietary Substances by Use of a Microrespirometer Technique: I. In Vitro Study of Kasdenol. Annals of Dentistry, December 1961.
- 28) Volpe, A. R., Manhold, J. H.: Microrespirometer Technique for Study of Human Gingival Tissue. Jour. Dent. Research 41: 420, 1962.
- 29) Manhold, J. H., Parker, L. A., and Manhold, B. S.: Efficacy of a Commercial Mouth Wash; In Vivo Study. N. Y. Jour. Dent. 32:165, 1962.
- 30) Manhold, J. H.: Gross Pathology, Histopathology and Psychopathology of Joint Disturbances. N. J. State Dent. Jour. May-June 1962.
- 31) Volpe, A. R., and Manhold, J. H.: A Method of Evaluating the Effectiveness of Potential Calculus Inhibiting Agents. N. Y. State Dental Jour. 28:289, 1962.
- 32) Volpe, A. R., Manhold, J. H., and Manhold, B. S.: Effect of Age and Other Factors Upon Normal Gingival Tissue Respiration. Jour. Dent. Research 41: 1060, 1962.
- 34) Manhold, J. H.: Comparison of Dental Caries Ratings: A Statistical Note. N. Y. Jour. Dent. 33:10, 1962.

1005101393

- 35) Manhold, J. H., Vinton, P. M., and Manhold, B. S.: Preliminary Notes on a Study of the Efficacy of the DAT in Predicting Four Year Performance in a New School. Jour. Dent. Education 27:84, 1963.
- 36) Manhold, J. H., and Volpe, A. R.: Effect of Inflammation in the Absence of Proliferation on  $O_2$  Consumption of Gingival Tissue. Jour. Dent. Research 42:103, 1963.
- 37) Manhold, J. H., Volpe, A. R., and DeCroce, R.: Relationship Between Salivary Respiration and Periodontal Disease: A Preliminary Report. Preprinted abstracts of the I.A.D.R. Meeting of March 1963.
- 38) Manhold, J. H., Volpe, A. R. and DeCroce, R.: Amino Acid Composition of Human Gingival Tissue as Determined by Paper Chromatography: A Preliminary Report. Preprinted abstracts of the I.A.D.R. Meeting of March 1963.
- 39) Manhold, J. H., Shatin, Leo, and Manhold, B. S.: A Comparison of Dental and Medical Students with Respect to interests, needs, and selected personality factors. Jour. Am. Dent. Asso. 67:601, 1963.
- 40) Manhold, J. H., Jr.: Oral Manifestations of Psychosomatic Disease. Psychosomatics Jour. IV, #5, Sept. - Oct., 1963, pg. 279.
- 41) Lapeyrolerie, F. M., and Manhold, J. H.: Hepatoma Metastatic to the Gingiva. Jour. Oral Surg., Oral Med., and Oral Path. 18:365, 1964.
- 42) Manhold, J. H., and Manhold, B. S.: Further in Vivo Study of Commercial Mouth Wash Efficacy. N. Y. Jour. Dent. 33:383, 1963.
- 43) Manhold, J. H., Manhold, B. S., and Volpe, A. R.: Relation Between Salivary Calculus and Blood Serum Phosphorus. Abstract, Jour. Dent. Research 43:838, 1964.
- 44) Volpe, A. R., Manhold, J. H., and Manhold, B. S.: Relationship Between Calculus Formation and Salivary Respiration ( $QO_2$ ): A Preliminary Report. Abstract, Jour. Dent. Research 43:838, 1964.
- 45) Mohammed, C. I., Manhold, J. H., and Manhold, B. S.: Efficacy of Pre-operative Oral Rinsing to Reduce Air Contamination During the Use of Air Turbine Handpieces. Jour. Am. Dent. Asso. 69:715, 1964.
- 46) Manhold, J. H.: Dental Psychosomatic Aspects of Pediatrics. Psychosomatics, V: 358, 1964.
- 47) Manhold, J. H. and Manhold, B. S.: Final Report of An Eight-Year Study of the Efficacy of the Dental Aptitude Test in Predicting Four Year Performance in a New Dental School. Jour. Dent. Ed. 29:41, 1965.
- 48) Volpe, A. R., Manhold, J. H., and Hazen, S. P.: In Vivo Calculus Assessment: Part I: A Method and its Examiner Reproducibility. Jour. Perio. 36:292, 1965.

1005101394

- 49) Manhold, J. H., Volpe, A. R., Hazen, S. P., Parker, L. A., and Adams, S. H.: In Vivo Calculus Assessment: Part II: A Comparison of Scoring Techniques. Jour. Perio. 36:299, 1965.
- 50) Manhold, J. H.: Dental Management of the Young Child and Old Adult. IN PRESS Proceedings of the 125th Anniversary of the College of Dentistry of the University of Maryland.
- 51) Manhold, J. H., Franzetti, J., and Fitzsimmons, L.: Effect of the Electric Toothbrush on Human Gingiva: Histologic and Microrespirometer Evaluation. Jour. Perio. 36:135, 1965.
- 52) Hazen, S. P., Volpe, A. R., and Manhold, J. H.: Relationship Between the Calculus Present on Teeth and Stainable Dental Plaque. Jour. Perio. 36:394, 1965.
- 53) Manhold, J. H., Volpe, A. R., and Manhold, B. S.: Inter-relationship Between Salivary Calculus Formation, Blood Serum Phosphorus Level, and Salivary Respiration ( $\text{CO}_2$ ). Jour. Perio. 36:474, 1965.
- 54) Manhold, J. H.: General Principles of Growth and Its Disturbances. Jour. N. J. State Dent. Soc. 37:25, 1965.

1005101395

PRELIMINARY REPORT OF RESPIRATION CHANGES IN RAT GLAND TISSUE AS A RESULT OF LIGATION. J.H. Manhold, Jr., T. E. Bolden, and M. L. Hall. The exorbital lacrimal gland of the rat exhibits many similarities to the salivary gland, and is more advantageous for study. A normal respiratory quotient ( $QO_2 = 0.41 \pm 0.14$ ) was established for exorbital lacrimal gland tissue of 150 day old male Wistar rats by use of a microrespirometer technique (Manhold, Bolden, and Katz 1959). Excretory ducts were then ligated, and the animals killed at 1, 2, 3, 7, 14 days, 1 month, and 4 months post-ligation. The duct-ligated glands were removed, placed in Tyrode's solution, and divided into two parts. Half was fixed for microscopic study. Half was placed in microrespirometers for  $QO_2$  determination. Results were: 1 day - (11 glands)  $0.36 \pm 0.59$ ; 2 days - (9 glands)  $0.34 \pm 0.22$ ; 3 days - (8 glands)  $0.70 \pm 0.08$ ; 7 days - (12 glands)  $0.64 \pm 0.53$ ; 14 days - (11 glands)  $0.50 \pm 0.14$ ; 1 month - (10 glands)  $0.29 \pm 0.25$ ; 4 months - (10 glands)  $0.30 \pm 0.19$ . These results corresponded with the microscopic picture. Tentatively, therefore, it may be concluded that duct-ligation caused an initial drop in the respiration of glandular tissue followed on the third day by an increase. A constant  $QO_2$  value, which is lower than the initial value, is reached when the gland histologically shows more scar tissue and less parenchyma.

Presented at the 39th General Meeting of the International Association for Dental Research, Boston, Massachusetts, March 1961.

#### COMPARATIVE STUDY OF SUBMAXILLARY AND LACRIMAL GLANDS OF THE RAT.

Theodore E. Bolden and John H. Manhold, Jr. Striking similarities in composition and response to disease processes have been demonstrated for salivary and lacrimal glands. This study determined the composition and dry weight of lacrimal glands and compared some of the data with data obtained in a similar fashion for submaxillary glands. Lacrimal glands from seven groups of untreated male Albino rats were used. (15, 30, 60, 90, 150, 300 and 500 days.) A quantitative estimate was made of the relative areas in sections of left glands which were occupied by acinar and by intralobular duct tissue. An ocular grid divided into 66 squares was used. A total of 1200 squares were counted. The number of squares in which half or more of the area was occupied by acini or ducts was counted in 3 specimen from each age group. Dry weight determinations were made on the right gland of 27 rats from the last six ages. The relative proportion of acinar tissue to intralobular duct tissue remained fairly constant at all ages. The normal gland consisted of about 82% acinar tissue, and 3% intralobular duct tissue. Lacrimal glands consisted of approximately 24% more acinar tissue and 22% less intralobular duct tissue than the submaxillary glands from rats of the same age. An average dry weight of 8 mg. (30 days) was increased to 50 mg. (60 days) fell, and fluctuated around 25 mg. at all other ages. The ratio of gland weight to body weight was 0.75 and .189 respectfully, for lacrimal and submaxillary glands.

Presented at the 39th General Meeting of the International Association for Dental Research, Boston, Massachusetts, March 1961.

1005101396

MICRORESPIROMETER TECHNIC FOR STUDY OF HUMAN  
GINGIVAL TISSUE: PRELIMINARY REPORT.

J. H. Manhold, Jr., T. E. Bolden, and S. Katz,  
Seton Hall College of Medicine and Dentistry,  
Jersey City.

Glickman, Turesky, and Hill (1949) employed the Warburg manometric technic to study human gingival metabolism. Because accurate Warburg determinations require relatively large quantities of tissue (40-50 mg. wet weight) and such quantities of human tissue usually are not readily available, Glickman, Turesky, and Manhold (1950) employed dogs to study oxygen requirements of healing gingival tissue. Exact relationships between human and dog gingiva  $Q_{O_2}$  values were not found. Further investigations did not follow. Kirk (1950) presented a quantitative ultramicroanalysis method for measuring the metabolism of tissue cultures. This technic, giving accuracy on 10 mg. or less (wet weight) of tissue, appeared ideal for human gingival study and was used in the present investigation. Small sections of gingival tissue, removed from clinic patients of Seton Hall College of Medicine and Dentistry, were placed in Tyrode's solution and divided into two approximately equal parts. Half was fixed for microscopic study. Half was placed in microrespirometers for  $Q_{O_2}$  determinations. Thirty-six samples of microscopically determined normal and essentially normal gingival tissue yielded a mean  $Q_{O_2}$  of  $1.39 \pm 0.46$ , a value comparable to earlier results. Microrespirometer investigation of human gingival tissue appears feasible, therefore, and presents the advantage of using small tissue quantities.

1005101397